

# Finding Unresolved Brown Dwarf and Exoplanet Companions to M Dwarfs Using Mid-infrared Excess

N. J. Cook<sup>1</sup>, D. J. Pinfield<sup>1</sup>, B. Burningham<sup>1</sup>, A. C. Day-Jones<sup>1</sup>,

<sup>1</sup>Centre for Astrophysics Research, Science and Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB

## Abstract

We present an optimised method for identifying brown dwarf and exoplanet companions to very low mass stars. We identify an all sky sample of bright M dwarfs based on optical and near-infrared colours, reduced proper motion, with strict E(H-W2) constraints and H-W2 photometric uncertainty less than 4%. We hunt for excess in the mid infrared using H-W1 and H-W2 colours, and comparison samples of other M dwarfs from common multi-colour parameter-space (not including H-W1 and H-W2). These candidates will be followed up with adaptive optics, radial velocities, and light curves (for transit) where appropriate.

## Introduction

Locating unresolved ultra-cool (brown dwarf and giant exoplanet) companions to M dwarfs is vital to understand the fundamental nature of these sub-stellar objects. The properties of ultra-cool dwarfs (UCDs) are strongly dependent on mass and age. Thick cloudy atmospheres mean that UCD properties also depend on effective temperature, surface gravity and metallicity. For a complete understanding of UCDs it is important to have accurate measurements of mass, radius, metallicity and luminosity data. It is for this reason that a UCD as a companion allows us to measure all of these properties. As UCDs are inherently faint, the use of faint M dwarfs as primaries are ideal to enable unresolved companions. With the recent full-sky release (March 2012) of the Wide-Field Infrared Survey Explorer, (WISE, Wright et al. 2010) and the Two Micron All Sky Survey (2MASS, Skrutskie et al. 2006), we present an optimised method for finding ultra-cool companions to M dwarfs based on identifying colour excesses. This method would not be possible without the vast number of sources present in the WISE and 2MASS databases.

## Method

We constructed an initial sample by cross matching WISE sources with 2MASS, using a set of basic cuts, including removing the galactic plane at a galactic latitude of  $\pm 15^\circ$ . We chose the colours H-W1 and H-W2 as the best candidates to determine excess. The cuts comprised of cuts in colour of V-J greater than 4, and the J, H, K colour cuts, as well as the reduced proper motion cut defined in Lépine & Gaidos (2011). We required un-reddened photometry with an extinction E(H-W2) no greater than 3%, equivalent to A(V) less than 0.22. Photometric quality cuts of 4% were applied for H-W1 and H-W2, as well as identifying variable sources (Pinfield et al. 2013) and requiring all proper motions had greater than 4σ uncertainties, giving 78,454 M dwarf candidates. We identified similar M dwarfs using a control sample and a small volume in the colour space, V-J, J-H, and H-K around each of the M dwarf targets (see figure 1). Targets with few to no sources in their sub volume (SV) were rejected leaving 68,349 candidates. The mean and standard deviation (for H-W1, H-W2 and H - 0.5(W1-W2)) of each target's colour space SV was recorded.

We calculated the excess by subtracting the H-W1 and H-W2 colours from the mean of each target SV assuming that M dwarfs with similar V-J, J-H, and H-K colours also have similar H-W1 and H-W2 colours. Using simulated M dwarf-brown dwarf systems, and a control sample constructed using catalogues from Gliese & Jahreiß (1991) and Stauffer et al. (2010) we selected a specific region of excess-spectral type space (see figure 2). To reduce the number of candidates we increased the quality cut required in H-W1 and H-W2 and visual inspected the remaining candidates.




Figure 1: Colour-colour contour diagram for J-H and H-Ks illustrating the populations of M dwarfs used in the multi-colour-space analysis (V-J, J-H and H-K). Boxes represent targets and their respective sub-volumes, for which means and standard deviations in H-W1 and H-W2 were recorded.

## Summary

With the recent full data release of WISE combined with 2MASS we have constructed a new sample of M dwarfs. Using colour, reduced proper motion and photometric quality selections we select 78,454 high quality M dwarf candidates that occupy un-reddened regions of the sky. These M dwarfs have 2MASS J band magnitudes between 10 and 14.5. We use mid-infrared excess to select 68,349 of these candidates identifying colour excess by comparing similar M dwarfs in multi-colour-space.

By simulating M dwarf-brown dwarf systems we select our final candidates, and expect to find companions through a full follow-up programme. Using our multi-colour-space approach, made possible with a large WISE-2MASS sample, reduces the standard deviations in colour excess by a factor of ~2, increasing the chances of possible candidates detections in a full follow-up program.




Figure 2: Contour plot to show our sample of M dwarfs after the colour space analysis. The three lines represent the positions of simulated M dwarf-brown dwarf systems run through the multi-colour-space analysis. The candidates were selected by choosing an area between simulated primaries with no overall excess (middle lines), and one standard deviation excess (top line) with a one standard deviation deficit for comparison (bottom line). All analysis was completed in excess per standard deviation space. The rectangle represents the region in which final candidates were selected subject to visual inspection.

■ M dwarfs sources N = 108498  
 — No Excess :  $M + L3 \rightarrow M + L9.0$   
 —  $\sigma$  Excess :  $M + L3 \rightarrow M + L9.0$   
 —  $-\sigma$  Excess :  $M + L3 \rightarrow M + L9.0$   
 ■ Candidate selection box

## References

- Gliese, W., Jahreiß, H. 1991, On: The Astronomical Data Center CD-ROM: Selected Astronomical Catalogs, Vol. 1; L.E.-Brotzmann, S.E.-Gesser (eds.), NASA/Astronomical Data Center, Goddard Space Flight Center, Greenbelt, MD.  
 Lépine, S., Gaidos, E. 2011, *Astronomical Journal*, 142, 138  
 Pinfield, D., Gomes, J., Day-Jones, A. 2013, *European Physical Journal Web of Conferences*, 47, 6004  
 Roeser, S., Demleitner, M., Schilbach, E. 2010, *Astronomical Journal*, 139, 2440  
 Skrutskie, M.-F., Cutri, R.-M., Stiening, R., et al. 2006, *Astronomical Journal*, 131, 1163  
 Stauffer, J., Tanner, A.-M., Bryden, G., et al. 2010, *Publications of the ASP*, 122, 885  
 Wright, E.-L., Eisenhardt, P.-R.-M., Mainzer, A.-K., et al. 2010, *Astronomical Journal*, 140, 1868

## Acknowledgements

This publication makes use of data products from the WISE, which is a joint project of the University of California, Los Angeles, and the JPL/CIT, funded by NASA, and 2MASS which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/CIT, funded by the NASA and the NSF. This research has made use of the NASA/IPAC Infrared Science Archive, which is operated by the JPL, CIT, under contract with NASA, the VizieR database catalogue access tool and SIMBAD database, operated at CDS, Strasbourg, France. This work is based in part on services provided by the GAVO data center and the data products from the Position and Proper Motion Extended-L, PPMXL database of Roeser et al. (2010). The work of NJC was supported by the STFC awarded to the University of Hertfordshire 2012.

## Contact:

Neil J Cook  
 Centre for Astrophysics Research,  
 Science and Technology Research Institute,  
 University of Hertfordshire, Hatfield AL10 9AB  
 Email: neil.james.cook@gmail.com, n.cook4@herts.ac.uk